

SAVING YOUR HIGH-VALUE ASH

A simple guide for homeowners and municipalities with true ash (*Fraxinus*) trees

Piera Siegert, NH State Entomologist



Emerald ash borer adult
Photo credit: N.W. Siegert

The problem: Emerald ash borer is an introduced and destructive pest of all North American true ash (*Fraxinus*) such as white, green, and black/brown ash. Trees infested with emerald ash borer will die from the infestation within 3-5 years unless treated. Management strategies to slow the spread of ash mortality can reduce *overall* emerald ash borer populations, but they may not save the ash tree in front of **your** house or in **your** park. Potential costs associated with emerald ash borer for municipalities and homeowners include:

- Costs to remove/replace/treat infested trees
- Loss of landscaping and community character
- Increased heating/cooling costs
- Reduction in property value
- Potential property damage/personal injury suits
- Increased power outages



Emerald ash borer killed tree in Concord, NH.
Photo credit: P.Y. Siegert



Ash-lined neighborhood in Lebanon, NH.
Photo credit: P.Y. Siegert

Although you cannot control the arrival of emerald ash borer on your property, you can decide what impact emerald ash borer will have by developing an emerald ash borer plan. This should be done regardless of proximity to known emerald ash borer populations. The first step is to stay informed about known emerald ash borer populations in the state (www.nhbugs.org or follow NH Bugs on facebook). Next, determine if you have ash trees, what size they are, where they are located, and if they add value to your property or community. Use local foresters and arborists, on-line calculators (www.extension.entm.purdue.edu/treecomputer/ and other sites) or smart phone apps (ARBORJETP for iphone and ipad and others) to estimate the costs associated with tree removal, replacement or treatment. Once you have determined your investment in ash and considered your budget, you can develop a plan for which trees will be removed, replaced or treated with insecticides when emerald ash borer arrives. Having a plan empowers you to make informed decisions about your property or community.


Don't let the beetle decide what to do with your trees!

More information about using insecticides for emerald ash borer on the reverse

WARNING: Insecticides are chemicals that can have health or environmental impacts. If you use insecticides always follow all label instructions or hire a licensed pesticide applicator.

impact note: Some studies have linked imidacloprid to CCD, a world-wide decline in honeybee populations. The body of evidence is inconclusive, but potential impacts to pollinators should be considered when initiating an insecticide regime. For information on insecticides used for EAB see [www.emeraldashborer.info/files/Potential Side Effects of EAB Insecticides FAQ.pdf](http://www.emeraldashborer.info/files/Potential_Side_Effects_of_EAB_Insecticides_FAQ.pdf).

Pollinator



Pesticides can be a useful tool for protecting valued ash trees. There are important considerations to keep in mind, however, when selecting an insecticide regimen:

- Proximity to generally infested area—insecticide treatment is only recommended in the red and orange areas (see map left or visit www.nhbugs.org for the most current information)
- Size of tree—measure the diameter (in inches) at 4.5 feet above the ground with a caliper or tape measure to get Diameter at Breast Height (DBH)
- Health of tree—systemic insecticides are less effective in trees that are already in decline
- Proximity to socially or environmentally sensitive habitats (like school properties, wells, or wetland areas)
- Mode of application of insecticide
- Effectiveness of treatment
- Cost of treatment
- Frequency of treatment

Not all emerald ash borer-approved insecticides are equally effective, nor are they all appropriate in every circumstance. Choosing an ineffective treatment for your conditions may result in product failure and is not cost-effective. Less effective treatments may prolong the life of an ash tree early in the invasion process but as neighboring untreated ash trees start showing signs of decline, indicating increasing local emerald ash borer populations, a more effective pesticide treatment may be necessary. There are resources available to help you assess the management options and products that are right for you. The table below summarizes recommended chemicals available for use. Active ingredients are listed, not trade names. Most formulations are only available for use by a licensed pesticide applicator. For more detailed information about available insecticides, visit:

www.emeraldashborer.info/files/Multistate_EAB_Insecticide_Fact_Sheet.pdf. More resources are also available through www.nhbugs.org and www.emeraldashborer.info, as well as by contacting an experienced certified arborist.

<i>See map above to determine your management zone.</i>		Ash is less than 18" DBH	Ash is greater than 18" DBH
Generally infested	Ash appear healthy	Imidacloprid, dinotefuran, or emamectin benzoate	Emamectin benzoate
	Ash are in decline	Emamectin benzoate	
	Ash are dead or with greater than 50% crown dieback	Tree removal. Insecticides unlikely to be effective.	
Expansion management zone		Imidacloprid, dinotefuran, or emamectin benzoate	Emamectin benzoate
Alert management zone		Treatment not yet warranted. Develop a plan.	

For assistance in developing a municipal emerald ash borer plan, and to find out how your management goals coincide with state management of emerald ash borer, please contact your local UNH Cooperative Extension County Forester.